arXiv:1510.00267v1 [math.AG] 1 Oct 2015

NON-EXISTENCE OF HYPERSURFACES WITH REAL FIBERED LOGARITHMIC GAUSS MAP

KRISTIN SHAW

Abstract. This paper considers real algebraic hypersurfaces in $(\mathbb{C}^*)^{n+1}$ which are torically non-singular and have real fibered logarithmic Gauss map. It is proved that the only such hypersurfaces are hyperplanes in projective space.

1. Introduction

Simple Harnack curves, first introduced by Mikhalkin in [9], are real algebraic curves in toric surfaces which are characterized by various equivalent maximality properties (see [9], [11]). One of these defining properties is that the logarithmic Gauß map of a simple Harnack curve is real fibered. The adjectives *real fibered* and *totally real* from [11], [3] are synonymous.

Definition 1.1. *A morphism $f : X \to Y$ of real varieties is real fibered if $f(x) \in \mathbb{R}Y$ if and only if $x \in \mathbb{R}X$, where $\mathbb{R}X, \mathbb{R}Y$ denote the real parts of X, Y respectively.*

There has been much interest in determining the higher dimensional analogues of simple Harnack curves [1]. One proposed topological definition was given in [10]. Another proposed extension to higher dimensions, has been to axiomatize the defining property of Harack curves in terms of the real fibered logarithmic Gauß map, (see [10], [3]). Requiring the logarithmic Gauß map to be real fibered has also been used to define Harnack curves in the singular situation [8].

In [3], it was shown that if the Newton polytope of X is a size d simplex and X is torically non-singular (see Definition 1.2), then the logarithmic Gauß map is real fibered if and only if $d = 1$. In fact, the argument in [3] uses a less restrictive condition; namely that the logarithmic Gauß map is only required to be generically real fibered. Here, restricting to the case when the logarithmic Gauß map is real fibered everywhere, the proof of non-existence of such hypersurfaces is simplified and also generalized to all Newton polytopes which are non-singular in dimension 1. The proof uses a result from [7] which says that a real fibered map is unramified over the real locus, (see Theorem 1.7).

For a hypersurface $X \subset (\mathbb{C}^*)^{n+1}$, the logarithmic Gauß map can be defined by first choosing a branch of the holomorphic logarithm and then

2000 *Mathematics Subject Classification.* Primary 14P25; Secondary 14P05.

This research is supported by an Alexander von Humboldt Foundation Postdoctoral Research Fellowship.

composing with the usual Gauß map. Alternatively, if $X \subset (\mathbb{C}^*)^{n+1}$ is a hypersurface defined by a polynomial $f(z_0, \ldots, z_n)$ then the logarithmic Gauß map is given explicitly by:

$$\gamma(z_0, \ldots, z_n) = [z_0 \frac{\partial f}{\partial z_0} : \cdots : z_n \frac{\partial f}{\partial z_n}],$$

see [5]. The degree of the log Gauß map $\gamma_X : X \to \mathbb{C}P^n$ is the normalized volume of the Newton polytope Δ of the polynomial defining X, i.e.

$$\deg(\gamma_X) = \mathrm{Vol}_{n+1}(\Delta). \tag{1.1}$$

The normalized volume of a $n+1$ dimensional lattice polytope $\Delta \subset \mathbb{R}^{n+1}$ is $(n+1)!$ times its Euclidean volume, (see Definition 3.1).

Consider the closure $\overline{X}$ of X in the toric variety $\mathbb{C}(\Delta)$ associated to the Newton polytope Δ of X.

Definition 1.2. *A hypersurface $X \subset (\mathbb{C}^*)^{n+1}$ with Newton polytope Δ is torically non-singular if its closure $\overline{X} \subset \mathbb{C}(\Delta)$ is non-singular and the intersection of $\overline{X}$ with each toric orbit of $\mathbb{C}(\Delta)$ is non-singular in that orbit.*

The fact that $\overline{X}$ is non-singular for a torically non-singular hypersurface $X \subset (\mathbb{C}^*)^{n+1}$ implies that the Newton polytope of X is non-singular in dimension 1, see [4, Chapter 2].

Definition 1.3. *An $n+1$ dimensional lattice polytope $\Delta \subset \mathbb{R}^{n+1}$ is non-singular in dimension 1 if for for every edge E of Δ the primitive normal vectors to the facets of Δ adjacent to E are a subset of a $\mathbb{Z}^{n+1}$ basis.*

The theorem to be proved here is the following.

Theorem 1.4. *A torically non-singular real hypersurface $X \subset (\mathbb{C}^*)^{n+1}$ has real fibered logarithmic Gauß map if and only if its Newton polytope $\Delta \subset \mathbb{Z}^{n+1}$ is up to integer affine transformation the standard simplex. In particular, $\overline{X}$ is a real hyperplane in $\mathbb{C}P^{n+1}$.*

To prove the theorem it is first noted that the logarithmic Gauß map can be extended continuously over the closure $\overline{X}$ and there is the following result about this extension.

Proposition 1.5. [3, Proposition 1.5] *If $X \subset (\mathbb{C}^*)^{n+1}$ is a torically non-singular real hypersurface which has real fibered logarithmic Gauß map then the extension $\overline{\gamma} : \overline{X} \to \mathbb{C}P^n$ is also real fibered.*

The next statement also follows immediately from the proof of the above proposition in [3].

Corollary 1.6. *If $X \subset (\mathbb{C}^*)^{n+1}$ be a torically non-singular hypersurface, then for every face F of Δ of dimension k, the logarithmic Gauß map of $\overline{X} \cap \mathbb{C}(F)$ considered as a hypersurface of $\mathbb{C}(F)$ coincides with the restriction of $\gamma_{\overline{X}}$.*

The following theorem about real fibered morphisms is used to restrict the topology of $\mathbb{R}\overline{X}$.

Theorem 1.7. [7, Theorem 2.19] *Let $\phi : X \to Y$ be a totally real fibered morphism between equidimensional non-singular real varieties, then $d_x\phi : T_x\mathbb{R}X \to T_{\phi(x)}\mathbb{R}Y$ is an isomorphism for all $x \in \mathbb{R}X$.*

Here the proof of the above theorem is outlined in the setting when $Y = \mathbb{C}P^n$ referring the reader to [7] for details. Firstly, notice that the statement is true when X and Y are 1-dimensional. To see this, it is enough to work locally; if a real map $\phi : (\mathbb{C}, 0) \to (\mathbb{C}, 0)$ is ramified at 0, then ϕ is locally given by $z \mapsto z^n$ for $n \geq 2$ which is clearly not real fibered.

Now for a map $\phi : X \to \mathbb{C}P^n$, where X is non-singular and of dimension n, it is enough to prove that $d_x\phi$ is injective. Suppose there is a point $x \in \mathbb{R}X$ such that $d_x\phi : T_x\mathbb{R}X \to T_{\phi(x)}\mathbb{R}P^2$ is not injective. Then choose a real line $L \subset \mathbb{C}P^n$ through $\phi(x)$ and such that $T_{\phi(x)}\mathbb{R}L \cap d_x\phi(T_x\mathbb{R}X) = 0$. Consider the preimage $C = \phi^{-1}(L) \subset X$, and its normalization $\pi : \tilde{C} \to C$. The composition $\phi \circ \pi : \tilde{C} \to L$ is also a real fibered map, see [7, Theorem 2.15, Corollary 2.16]. Since the theorem is true for maps between curves, this map is unramified over the real locus.

However, for any point $\tilde{x} \in \mathbb{R}\tilde{C}$ such that $\pi(\tilde{x}) = x \in \mathbb{R}C \subset \mathbb{R}X$, the differential satisfies $d_{\tilde{x}}(\phi \circ \pi) = d_x\phi \circ d_{\tilde{x}}\pi$. Therefore the image of $d_{\tilde{x}}(\phi \circ \pi)$ is zero by the assumption that $T_{\phi(x)}\mathbb{R}L \cap d_x\phi(T_x\mathbb{R}X) = 0$. This gives a contradiction and the theorem follows.

If the logarithmic Gauß map $\gamma : \overline{X} \to \mathbb{C}P^n$ is real fibered, and the closure $\overline{X}$ in $\mathbb{C}(\Delta)$ is non-singular, then it follows that $\gamma : \mathbb{R}\overline{X} \to \mathbb{R}P^n$ is a covering map, see [7, Corollary 2.21]. For $n > 1$, fundamental group of $\mathbb{R}P^n$ is $\mathbb{Z}/\mathbb{Z}_2$. So there are only two connected coverings; $\mathbb{R}P^n \to \mathbb{R}P^n$ of degree 1 and $S^n \to \mathbb{R}P^n$ of degree 2. In [7], a real projective variety X of dimension n admitting a real fibered morphism to $\mathbb{C}P^n$ is called *weakly hyperbolic*. For $n = 1$, weak hyperbolicity is equivalent to being a curve of dividing type, also known as type one [7, Theorem 2.9].

For $n > 1$, if a variety is weakly hyperbolic, the degree of a map to $\mathbb{C}P^n$ is determined by the topology of $\mathbb{R}X$. Applying Formula 1.1 for the degree of the logarithmic Gauß map gives the following.

Corollary 1.8. *For $n > 1$, it $X \subset (\mathbb{C}^*)^{n+1}$ is a torically non-singular real hypersurface with real fibered logarithmic Gauß map then $\mathbb{R}\overline{X}$ is a disjoint union of k copies of S^n and l copies of $\mathbb{R}P^n$ where*

$$\deg(\gamma) = Vol_{n+1}(\Delta) = 2k + l.$$

To prove Theorem 1.4, Section 2 first considers the case of surfaces $X \subset (\mathbb{C}^*)^3$ with real fibered logarithmic Gauß map. By Proposition 1.5, the intersection of the closure $\overline{X}$ with a toric invariant divisor of $\mathbb{C}(\Delta)$ is a simple Harnack curve in the toric surface $\mathbb{C}(F)$ for each facet F of Δ. Using the topological description of Harnack curves and simple consequences of Theorem 1.7, it is proved in Proposition 2.2, that the Newton polytope of a surface with real fibered logarithmic Gauß map must be a lattice tetrahedron whose edge lengths are all 1.

Finally, in Section 3, Theorem 1.4 is proved in full generality by applying an induction argument on n with $n = 2$ being the base case. The proof uses Lemma 3.3 which shows that a lattice simplex, non-singular
in dimension 1, with unimodular facets (see Definition 3.1) is up to integral affine transformation the standard simplex.

The next example provides a counter-example to Theorem 1.4 if the assumption that X be torically non-singular is dropped.

Example 1.9. Requiring that the closure $\overline{X}$ of X in $\mathbb{C}(\Delta)$ be non-singular is necessary for Theorem 1.4. For example, let $\Delta \subset \mathbb{R}^3$ be the convex hull of

$$(0,0,0),(1,0,0),(0,1,0),(0,0,2).$$

At the edge $[(1,0,0),(0,1,0)]$, the polytope Δ fails the smoothness condition, since the normals of the adjacent facets are $(0,0,-1)$ and $(2,2,1)$, which are not a subset of a $\mathbb{Z}^3$ basis.

However, the hypersurface $X \subset (\mathbb{C}^*)^3$ defined by the equation

$$f(z_1,z_2,z_3) = z_3^2 + az_3 + bz_1 + cz_2 + d,$$

with $a,b,c,d \in \mathbb{R}$ satisfying $a^2 - 4d > 0$ has totally real fibered logarithmic Gauß map $\gamma : X \to \mathbb{C}P^2$.

Although $\overline{X}$ is singular, one can consider the desingularization $\mathbb{C}(\tilde{\Delta})$ of $\mathbb{C}(\Delta)$ and let $\tilde{X}$ be the closure of X in $\mathbb{C}(\tilde{\Delta})$. Then the logarithmic Gauß map $\gamma : \tilde{X} \to \mathbb{C}P^2$ is real fibered outside of a point in $\mathbb{C}P^2$, but not real fibered everywhere.

Acknowledgements: The author is grateful to Erwan Brugallé, Christian Haase and Mario Kummer for helpful discussions. In particular to Erwan Brugallé for pointing out Example 1.9. Thanks also to Michael Joswig for providing helpful comments on a preliminary draft.

2. The surface case and boundary Harnack curves

Suppose $X \subset (\mathbb{C}^*)^3$ is a torically non-singular real surface with real fibered logarithmic Gauß map. By Proposition 1.5, for each facet F of Δ each curve $Z = \overline{X} \cap \mathbb{C}(F)$ has real fibered logarithmic Gauß map, hence it is a simple Harnack curve. By the topological characterization of Harnack curves (see [9]), the real part $\mathbb{R}Z$ has a single connected component intersecting the toric boundary divisors of the surface $\mathbb{C}(F)$. Call this connected component the *outer connected component* of the curve Z and denote it by $O(Z) \subset \mathbb{R}Z$.

There is the following formula for the degree of the logarithmic Gauß map restricted to $O(Z)$ in terms of the lattice perimeter of the face F,

$$\deg(\gamma|_{O(Z)}) = -2 + \sum_{E \in \mathcal{E}(F)} \text{Length}(E), \tag{2.1}$$

where $\text{Length}(E) := |E \cap \mathbb{Z}^n| - 1$, see [11].

Moreover, Harnack curves have the property that all intersections with toric invariant curves of the surface $\mathbb{C}(F)$ are real and transverse. That X is torically non-singular implies for two facets F and F' of Δ the corresponding outer components $O(Z)$ and $O(Z')$ also intersect transversally and in exactly $\text{Length}(E)$ points where $E = F \cap F'$.

Lemma 2.1. *If $X \subset (\mathbb{C}^*)^3$ is a torically non-singular real surface with real fibered logarithmic Gauß map, then there exists a connected component $\mathcal{C} \subset \mathbb{R}\overline{X}$ containing the outer components of all the toric invariant boundary curves of $\overline{X}$.*

Proof. Let Δ be the Newton polytope of X. Each of the components $O(Z_i)$ is an embedded S^1 contained in some connected component of $\mathbb{R}\overline{X}$. Faces F and F' of Δ intersect in an edge E if and only if the corresponding components $O(Z)$ and $O(Z')$ intersect transversally in exactly Length(E) points. This means that $O(Z)$ and $O(Z')$ are contained in the same connected component of $\mathbb{R}\overline{X}$. The facets of Δ are connected via the edges therefore there exists a single component $\mathbb{R}\overline{X}$ containing the outer components of all boundary curves. □

Proposition 2.2. *If $X \subset (\mathbb{C}^*)^3$ is a torically non-singular real surface with real fibered logarithmic Gauß map, then the connected component $\mathcal{C} \subset \mathbb{R}(\overline{X})$ which contains the outer components of all of the simple Harnack boundary curves is homeomorphic to $\mathbb{R}P^2$.*

Moreover, the Newton polytope of X is a tetrahedron with all edges of lattice length 1*.*

Proof. By Corollary 1.8, the connected component $\mathcal{C}$ is homeomorphic to either S^2 or $\mathbb{R}P^2$. Suppose the connected component $\mathcal{C}$ containing all of the outer components is homeomorphic to S^2. Any two outer components $O(Z_i)$, $O(Z_j)$ intersect transversally in exactly Length(E) points in S^2. However two closed curves in S^2 intersecting transversally do so in an even number of points. This implies that Length$(E) = 2k > 0$ for each edge E of Δ. A facet F of Δ has at least 3 edges, therefore the normalized perimeter of every facet F satisfies $\sum_{E\in\mathcal{E}(F)} \text{Length}(E) \geq 6$.

On the other hand, the restriction of the logarithmic Gauß map $\gamma_{\overline{X}}$ to an outer component $O(Z)$ must have degree either 1 or 2, depending if the image $\gamma(O(Z))$ is an *oval* or a *pseudoline* respectively. Recall that an oval in $\mathbb{R}P^2$ is an embedded S^1 with trivial $\mathbb{Z}/2\mathbb{Z}$ homology class, and a pseudoline is an embedded S^1 realizing the generator of $H_1(\mathbb{R}P^2, \mathbb{Z}/2\mathbb{Z})$. So by Equation 2.1

$$\deg(\gamma|_{O(Z)}) = -2 + \sum_{E\in\mathcal{E}(F)} \text{Length}(E) = 1 \text{ or } 2.$$

Therefore, $\sum_{E\in\mathcal{E}(F_i)} \text{Length}(E) = 3$ or 4 which yields a contradiction to the lower bound of the lattice perimeter of F given above.

So $\mathcal{C}$ is homeomorphic to $\mathbb{R}P^2$ and the restriction of the logarithmic Gauß map $\gamma : \mathcal{C} \to \mathbb{R}P^2$ is 1-1. Using Equation 2.1 again gives,

$$\deg(\gamma|_{O(Z)}) = \sum_{E\in\mathcal{E}(F)} \text{Length}(E) - 2 = 1,$$

which implies that each face F is a lattice triangle and Length$(E) = 1$ for all edges E of Δ.

The outer component of each boundary curve must be a pseudoline, since if it was an oval, for any embedded circle Y in $\mathbb{R}P^2$ intersecting $O(Z)$ transversally, we have $|O(Z) \cap Y| = 2k$ for $k > 0$. However, there is a facet F' intersecting F in an edge of Δ. By above this edge has length 1, so that $O(Z)$ and $O(Z')$ intersect transversally in exactly 1 point, which gives a contradiction to $O(Z)$ being an oval.

Since each outer component is a pseudoline, for all pairs of faces F and F' the intersection $O(Z) \cap O(Z') \neq \emptyset$. Therefore, each pair of faces of Δ must

share an edge and there are exactly 4 faces of dimension 2. This implies that Δ is a lattice tetrahedron. Each edge of Δ was already shown to have lattice length 1, and the proposition is proved. □

The next corollary follows immediately from the previous proposition. Notice that this is also a direct consequence of the stronger result proved in [3], where it is only assumed that the logarithmic Gauß map be generically real fibered.

Corollary 2.3. *If $X \subset (\mathbb{C}^*)^3$ is a hypersurface with real fibered logarithmic Gauß map and Newton polytope a size d standard simplex then $d = 1$.*

3. Proof of main theorem

Let F be a face of dimension k of a lattice polytope $\Delta \subset \mathbb{R}^{n+1}$. Denote by V_F the k dimensional vector subspace of $\mathbb{R}^{n+1}$ parallel to F and let $\Lambda_F = V_F \cap \mathbb{Z}^{n+1}$ be the rank k lattice. Choose a basis $v_1, \dots, v_k$ of Λ_F and let $\mathcal{S}(\Lambda_F)$ be the k-dimensional lattice simplex which is the convex hull of $\{0, v_1, \dots, v_k\}$.

Definition 3.1. *The normalized volume of a k dimensional lattice polytope $F \subset \mathbb{R}^{n+1}$ is*

$$\mathrm{Vol}_k(F) = \frac{\mathrm{Vol}_k^E(F)}{\mathrm{Vol}_k^E(\mathcal{S}(\Lambda_F))},$$

where Vol_k^E denotes the Euclidean volume. Call a k-dimensional face $F \subset \Delta \subset \mathbb{R}^{n+1}$ unimodular if $\mathrm{Vol}_k(F) = 1$.

Remark 3.2. *The above definition does not depend on the choice of basis $v_1, \dots, v_k$ of Λ_F. Also notice that F is unimodular if and only if, up to integer affine transformation, it is the convex hull of $\{0, e_1, \dots, e_k\}$.*

Notice that for a lattice segment, $\mathrm{Length}(E) = \mathrm{Vol}_1(E)$. *Also denote* $\mathrm{Area}(F) := \mathrm{Vol}_2(F)$ *for a polygon $F \subset \mathbb{R}^{n+1}$*

Finally, if $\Delta \subset \mathbb{R}^{n+1}$ is an $n+1$ dimensional lattice simplex with vertices $0, v_1, \dots, v_{n+1}$ then $\mathrm{Vol}_{n+1}(\Delta) = |\det(v_1, \dots, v_{n+1})|$.

Lemma 3.3. *If $\Delta \subset \mathbb{R}^{n+1}$ is an $n+1$ dimensional lattice simplex with unimodular facets, which is also smooth in dimension 1, then Δ is unimodular.*

Proof. Choose a facet F of Δ. Since F is unimodular, it can be assumed, up to an integer affine transformation, that the vertices of F are given by $\{0, e_1, \dots, e_n\}$. There is one additional vertex of Δ, to be denoted by a where $a = (a_1, \dots, a_n, v) \in \mathbb{Z}^{n+1}$. Note that $\mathrm{Vol}(\Delta) = |\det(e_1, \dots, e_n, a)| = v$.

By assumption, the facet with vertices $\{0, e_1, \dots, \hat{e_i}, \dots, e_{n-1}, a\}$ is also unimodular, so there is a vector $c = (c_1, \dots, c_{n+1}) \in \mathbb{Z}^{n+1}$ such that

$$\det(c, e_1, \dots, \hat{e_i}, \dots, e_{n-1}, a) = \pm(vc_i \pm a_i c_n) = \pm 1.$$

Therefore, $\gcd(v, a_i) = 1$ and the primitive normal vector to this facet is $\pm(a_i e_{n+1} \pm v e_i)$.

The condition that Δ is smooth in dimension 1 implies that at each edge of Δ the normal vectors of the facets adjacent to that edge are a subset of a $\mathbb{Z}^{n+1}$ basis. Apply this condition at the edge $[0, e_1]$ of Δ, where the normal

vectors of the adjacent facets are $a_i e_{n+1} - v e_i$ for $i = 2, \dots n$ and e_{n+1}. Once again there must exist a $c = (c_1, \dots, c_{n+1}) \in \mathbb{Z}^{n+1}$, such that

$$\det(c, a_2 e_{n+1} - v e_2, \dots, a_n e_{n+1} - v e_n, e_{n+1}) = \pm c_1 \cdot v^{n-1} = \pm 1.$$

Therefore, $\mathrm{Vol}(\Delta) = v = 1$ and Δ is unimodular and the lemma is proved. □

Remark 3.4. *Notice that both assumptions are necessary for the above lemma to hold. Already in dimension* 3 *there are tetrahedra, with unimodular faces which are not unimodular, for example the convex hull of*

$$(0,0,0), (1,0,0), (0,1,0), (1,p,q),$$

for every pair of p, q *with* $\gcd(p,q) = 1$*, has unimodular facets but the volume of the tetrahedron is* q*.*

The condition that the facets of Δ *be unimodular is also necessary, consider the trivial example of the size* d*-standard simplex.*

Proof of Theorem 1.4. The theorem is proved by induction on n, starting with $n = 2$ as the base case. By Proposition 2.2, the connected component $\mathcal{C}$ of $\mathbb{R}\overline{X}$ containing all the outer components must be $\mathbb{R}P^2$ and the Newton polytope Δ of X is a tetrahedron with all edge lengths equal to 1, so that,

$$\sum_{E \in \mathcal{E}(\Delta)} \mathrm{Length}(E) = 6, \qquad |\mathcal{F}(\Delta)| \geq 4, \qquad \text{and} \qquad l \geq 1, \tag{3.1}$$

where l is the number of connected components of $\mathbb{R}X$ homemorphic to $\mathbb{R}P^2$.

Let $\beta_*(X; \mathbb{Z}/2\mathbb{Z})$ denote the sum of all $\mathbb{Z}/2\mathbb{Z}$ Betti numbers. There is the Smith-Thom inequality (see [2]):

$$\beta_*(\mathbb{R}\overline{X}; \mathbb{Z}/2\mathbb{Z}) \leq \beta_*(\overline{X}; \mathbb{Z}/2\mathbb{Z}). \tag{3.2}$$

By Corollary 1.8, the real part $\mathbb{R}\overline{X}$ is a union of k disjoint 2-spheres and l real projective planes. The total sum of Betti numbers for $\mathbb{R}P^2$ and S^2 are,

$$\beta_*(\mathbb{R}P^2; \mathbb{Z}/2\mathbb{Z}) = 3 \qquad \text{and} \qquad \beta_*(S^2; \mathbb{Z}/2\mathbb{Z}) = 2.$$

So that

$$\beta_*(\mathbb{R}\overline{X}) = 3l + 2k = 2l + \mathrm{Vol}(\Delta). \tag{3.3}$$

Moreover, Khovanskii's formula [6] for the Euler characteristic of the complex hypersurface $\overline{X}$, gives,

$$\beta_*(\overline{X}; \mathbb{Z}/2\mathbb{Z}) = \mathrm{Vol}_3(\Delta) - \sum_{F \in \mathcal{F}(\Delta)} \mathrm{Area}(F) + \sum_{E \in \mathcal{E}(\Delta)} \mathrm{Length}(E). \tag{3.4}$$

Combining Equations 3.1, 3.3, 3.4 and with the Smith-Thom inequality 3.2 yields,

$$2l + \mathrm{Vol}_3(\Delta) \leq \mathrm{Vol}_3(\Delta) - \sum_{F \in \mathcal{F}(\Delta)} \mathrm{Area}(F) + \sum_{E \in \mathcal{E}(\Delta)} \mathrm{Length}(E).$$

Applying (3.1), further implies

$$\sum_{F \in \mathcal{F}(\Delta)} \mathrm{Area}(F) \leq 4.$$

Therefore, each facet of Δ is unimodular. Since Δ is also non-singular in dimension 1, Lemma 3.3 implies that Δ is unimodular and therefore $\mathbb{C}(\Delta)$ is $\mathbb{C}P^3$ and $\overline{X}$ is a hyperplane.

Now proceed by induction for $n \geq 3$. Suppose $X \subset (\mathbb{C}^*)^{n+1}$ is torically non-singular and has real fibered logarithmic Gauß map. Then by Proposition 1.5, for each toric invariant divisor $\mathbb{C}(F)$ of $\mathbb{C}(\Delta)$ corresponding to the facet F of Δ, the intersection $\overline{X} \cap \mathbb{C}(F)$ has real fibered logarithmic Gauß map. By applying the induction hypothesis, each facet F of Δ is unimodular and the toric variety $\mathbb{C}(F)$ is $\mathbb{C}P^n$, and $\overline{X} \cap \mathbb{C}(F)$ is a hyperplane for each facet F of Δ. In particular the real part, $\mathbb{R}\overline{X} \cap \mathbb{C}(F)$ is connected for all facets F.

Therefore, similar to Lemma 2.1, there is a single connected component $\mathcal{C}$ of $\mathbb{R}\overline{X}$ which contains all of the real subvarieties $\mathbb{R}\overline{X} \cap \mathbb{C}(F)$ for all faces F of Δ. Moreover, $\mathcal{C}$ is homeomorphic to $\mathbb{R}P^n$ since there is no embedding of $\mathbb{R}P^{n-1}$ to S^n for $n > 2$.

For each facet F of Δ, the class $[\mathbb{R}\overline{X} \cap \mathbb{C}(F)]$ is non-trivial in

$$H_{n-1}(\mathcal{C}; \mathbb{Z}/2\mathbb{Z}) = H_{n-1}(\mathbb{R}P^n; \mathbb{Z}/2\mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z}.$$

This is because for a 2 dimensional face A of Δ such that $A \cap F$ is an edge E, the varieties $\mathbb{R}\overline{X} \cap \mathbb{C}(F)$ and $\mathbb{R}\overline{X} \cap \mathbb{C}(A)$ intersect transversally, and their intersection is $\mathbb{R}\overline{X} \cap \mathbb{C}(E)$ which is a single point by the unimodularity of F. In particular, for any two facets, F, F' of Δ, the cup product of the Poincaré duals of $[\mathbb{R}\overline{X} \cap \mathbb{C}(F)]$ and $[\mathbb{R}\overline{X} \cap \mathbb{C}(F')]$ is a non-trivial cohomology class in $H^2(\mathbb{R}P^n; \mathbb{Z}/2\mathbb{Z})$. However, the intersection of $\mathbb{R}\overline{X} \cap \mathbb{C}(F)$ and $\mathbb{R}\overline{X} \cap \mathbb{C}(F')$ is of codimension 2 if and only if F and F' intersect in a face of Δ of codimension 2. This implies that every pair of facets of Δ must meet in a codimension 2 face. Therefore, Δ has at most $n+1$ facets, all of which are n dimensional unimodular lattice simplicies. Since Δ is also smooth in dimension 1 by assumption, applying Lemma 3.3 completes the proof. □

References

[1] Extreme forms of real algebraic varieties. Am. Inst. of Math. workshop, April 6th–9th 2006.

[2] R. Benedetti and J. J. Risler. *Real algebraic and semi-algebraic sets*: Actualités Mathématiques. [Current Mathematical Topics]. Hermann, Paris, 1990

[3] E. Brugallé, G. Mikhalkin, J-J. Risler. *Non-existence of torically maximal hypersurfaces*: arXiv preprint: arXiv:1506.02813.

[4] W. Fulton. *Introduction to toric varieties*, Princeton University Press, Princeton, NJ, 1993.

[5] M. Kapranov. *A characterization of A-discriminantal hypersurfaces in terms of the logarithmic Gauss map*, Math. Ann., **290(2)** (1991) 277–285.

[6] A. Khovanskii. *Newton polyhedra, and the genus of complete intersections*: Funktsional. Anal. i Prilozhen., **12(1)** (1978) 51–61.

[7] M. Kummer, E. Shamovich: *Real Fibered Morphisms and Ulrich Sheaves*, arXiv preprint: arXiv:1507.06760

[8] L. Lang: *A generalization of simple Harnack curves*, arXiv preprint: arXiv:1504.07256

[9] G. Mikhalkin: *Real algebraic curves, the moment map and amoebas*, Ann. of Math. **(2) , 151(1)** (2000), 309–326.

[10] G. Mikhalkin: *Amoebas of algebraic varieties*, arXiv preprint: arXiv:math/0108225, 2001.

[11] M. Passare, J-J. Risler: *On the curvature of the real amoeba*, In Proceedings of the Gökova Geometry-Topology Conference 2010, (2011),129–134.

Kristin Shaw, Technische Universität Berlin, MA 6-2, 10623 Berlin, Germany.

E-mail address: shaw@math.tu-berlin.de